



An exposomic framework to uncover environmental drivers of aging

Vrinda Kalia , PhD^{1,*}, Daniel W. Belsky , PhD², Andrea A. Baccarelli, MD, PhD¹, and Gary W. Miller , PhD¹

¹Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, NY 10032, USA

²Department of Epidemiology and Robert N. Butler Columbia Aging Center, Mailman School of Public Health, Columbia University, New York, NY 10032, USA

*To whom correspondence should be addressed: Email: vk2316@cumc.columbia.edu

Abstract

The exposome, the environmental complement of the genome, is an omics level characterization of an individual's exposures. There is growing interest in uncovering the role of the environment in human health using an exposomic framework that provides a systematic and unbiased analysis of the non-genetic drivers of health and disease. Many environmental toxicants are associated with molecular hallmarks of aging. An exposomic framework has potential to advance understanding of these associations and how modifications to the environment can promote healthy aging in the population. However, few studies have used this framework to study biological aging. We provide an overview of approaches and challenges in using an exposomic framework to investigate environmental drivers of aging. While capturing exposures over a life course is a daunting and expensive task, the use of historical data can be a practical way to approach this research.

Keywords: exposome; aging; hallmarks of aging; population-based studies of aging; measuring the exposome

Introduction

The human body reacts constantly to changes in its environment. Microbial invasions induce immunological responses [1]; social threats trigger the neuroendocrine cascade of the stress response [2] and can alter immune stance [3]; toxin, toxicant, and pollutant exposures elicit protective responses across multiple systems [4] and can induce adverse effects [5]; and change in ambient temperature can disrupt homeostasis [6]. On a cellular level, actions are taken every second based on signals received from the environment. Several factors can influence the effect of these exposures and the body's response, importantly, the levels of certain nutrients, and the overall nutritional status [7]. Environmental signals driving changes in our biology are constant while being variable in space and time [8]. Traditionally, scientists have studied the influence of the environment on human health using a reductionist approach, exploring the effect of a single environmental exposure on a health outcome. However, the recent emergence of the exposome concept provides an alternative framework to holistically study the environment. The exposome encompasses all exposures, throughout the life course [9], and includes internal processes, like endogenous metabolism and microbial-derived metabolites, specific external exposure, like exposure to toxic environmental chemicals and dietary nutrients, and wider social factors that can influence an individual, like financial status and education [10]. It represents the environmental complement to the genome, that is, a comprehensive characterization of the molecular exposures to which an organism is subject [11-13]. For the purpose of this review, we define the exposome as all exposures, including lived experiences, that can be assessed throughout the life course, but recognize

the need to focus on those exposures that can be measured with current technologies. We focus primarily on toxic exposures but we acknowledge other important factors that can influence the exposome or the body's response to exposure, such as: (i) nutritional status [7] and the microbiome [14] and (ii) other factors of the external exposome such as education [15] and family environment [16]. While there have been some attempts at characterizing the biological footprint of these factors [17], they remain difficult to measure. Newer studies provide hope and directions¹ on how these factors may be tested in the future [18].

The many chronic diseases of aging are thought to derive from common underlying processes that precipitate molecular changes over time [19]. These common mechanisms have been classified into nine hallmarks of aging: genomic instability, epigenetic alterations, telomere attrition, loss of proteostasis, deregulated nutrient sensing, mitochondrial dysfunction, cellular senescence, stem cell exhaustion, and altered intercellular communication [20]. Aging is influenced by genetic factors [21]; however, current estimates are that <10% of differences in longevity between individuals can be attributed to inherited genes [22]. Environmental factors are therefore critical determinants of aging processes [23].

The exposome and aging

The human body can activate compensatory pathways that provide a form of resilience against damaging factors that accumulate with age. It is proposed that individual differences in this process of resilience determine whether a life course would end relatively healthy, frail, or in early mortality [24, 25]. Adverse

Received: September 27, 2021. Revised: January 19, 2022. Accepted: January 24, 2022

© The Author(s) 2022. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

environmental exposures can induce damage and impose additional stress on resilience processes, reducing the reserve needed to counter the effect of biological aging, thus, increasing the risk of poor or accelerated aging.

Time is an important variable that can influence the effect of environmental exposures. Both the timing of exposure during a life course as well as the duration of an exposure can determine the physiological response. It is important to identify critical time windows of exposure across the life course that might influence an outcome of interest since the scale of the response to an exposure would depend on whether the exposure occurred in utero, during development, in mid-life, or later in life [26, 27]. Damaging environmental exposures may be for a short period of time (an acute exposure) or over a prolonged period (a chronic exposure). The response to an acute or transient exposure may subside once the exposure has been removed from the system through metabolism and excretion but may leave signs of lasting damage that can be detected through molecular probes, like epigenetic marks of exposure to cigarette smoke in former smokers [28] or epigenetic changes in immune cells in response to pathogenic exposure [29] whereas response to a chronic exposure is usually maintained to counter the effects of long-term exposure. Thus, exposures may induce temporary or permanent changes in molecular pathways. Both aspects of time can influence whether damage is caused and the organism's response to the exposure.

Several studies have reported associations between environmental exposures and different hallmarks of aging. For example, exposure to air pollution [30], pesticides [31], heavy metals [32], industrial solvents [33], and viruses and bacteria [34] has been associated with mitochondrial dysfunction but no study has investigated the effect of their concomitant exposure on mitochondrial function. Different environmental exposures are correlated, for example, people who live in polluted places tend to lead stressful lives and have reduced access to healthy nutrition, exercise, and leisure. For this reason, studying any one exposure may (a) confuse the apparent effects of the specific exposure with some other, correlated exposures and (b) fail to capture the myriad of exposures that represent the actual environmental burden. In order to illustrate the contributions of various exposures to hallmarks of aging, we have organized observed associations between exposures and the hallmarks of aging (Figure 1). This was created by first categorizing exposures into nine classes: air pollution, pesticides, metals/metalloids, mutagens, industrial solvents, plasticizers and related compounds, ambient temperature, microbes, and cigarette smoke. Each of these categories can comprise of multiple exposures and are grouped due to similarity in source, or chemical structure or properties. In order to confirm an association between the exposure class and a hallmark of aging, we ran multiple searches in PubMed using a combination of the name of each chemical category and, "aging," "hallmarks of aging," or "toxicity." The figure illustrates that different environmental exposures affect many different hallmarks of aging; therefore, an exposomic framework is needed to study environmental drivers of aging. Additionally, the influence of the exposome on the aging process would be modified by the underlying nutritional status and needs to be studied under these contexts [7, 35]. To the best of our knowledge, studies of aging have rarely adopted an exposomic framework.

Measuring the exposome

Compiling exposomic data requires integrating information from multiple sources. Broadly, the sources of data can be geospatial in origin or from personal monitoring [11, 36, 37].

"Geospatial data" can be used to infer individual-level exposure from ecological data. Using data describing the spatial distribution of a toxicant, an estimate of an individual's exposure to that toxicant can be made using their residential address. These data have been used to estimate exposure to air pollution, pesticides [38], access to green space, proximity to pollution sources, proximity to a contaminated water source, and proximity to landfills [39–41]. Data needed to measure the spatial distribution of a toxicant come from:

- a) Satellites in Earth's orbit which provide valuable information about the Earth's land surface, oceans, and atmosphere. Using optical density, radiation, and imagery data generated through satellite technology, researchers have been able to provide estimates of exposure to pollutants in the atmosphere, humidity, temperature, light exposure at night, even exposure to wildfires [42, 43]. Significant advances in the use of satellite data continue to be made, providing a means to get highly granular historical exposure data [44].
- b) *Ground-based monitors and sensors* distributed through national and regional networks have traditionally been used to monitor levels of various air pollutants and provide valuable data that can be used to attribute exposure to study participants using kriging methods [18, 19, 45]. The introduction of low-cost and mobile sensors has made it possible to add more nodes to the network of monitors and provide more granular estimates of air quality [46, 47].

"Personal monitoring data" provide individual-level exposure assessment that can be made through several methods:

- a) *Records and surveys* like the national health and nutrition examination survey, census surveys, records maintained by social services and criminal justice administrations, or personal records made through population studies can provide valuable information related to diet, lived experiences, adverse childhood experiences, and psychosocial stress. Census data and other administrative survey data can also be used to determine neighborhood level characteristics, like the racial/ethnic composition of a neighborhood [48, 49]. Personal health records, like electronic health records (EHRs), also provide a means to estimate exposures on a personal level, such as exposure to pharmaceutical drugs, recreational drugs, alcohol, or cigarette smoke. EHR also provide the aging relevant outcome data [50].
- b) *Wearable devices and smartphones* have made it possible to make personalized estimates of exposure. Data collected through accelerometers, GPS devices, and wearable sensors provide better estimates of individual level exposure [8, 51, 52]. This data can overcome issues related to geo-spatial data that do not account for movement and activity outside of the residence; however, most methods still need rigorous validation [53].
- c) *Biological samples* like blood, urine, sweat, saliva, feces, hair samples, or toenails provide a means to measure the burden of chemical exposures in an individual as well as the biological response as the result of an exposure. Improvements in high-resolution mass spectrometry instruments have made it possible to capture a large proportion of persistent pollutants in a tissue sample using appropriate chromatography [54]. Rapid progress in chemoinformatic software is improving confidence in annotations and expanding the known chemical space [55–57]. The relative levels of different chemicals in a biological matrix are a function of the time since




Figure 1. A selection of environmental exposures and hallmarks of aging. The nine hallmarks of aging represent common mechanisms of biological aging in the mammalian context. The inner most gray circle illustrates these hallmarks of aging. In the outer circle, icons represent environmental exposures that have been associated with each corresponding hallmark of aging. Each category can comprise of multiple exposures and are grouped due to similarity in: source, use, chemical structure or properties, or associated health effects. Created with [BioRender.com](https://www.biorender.com)

exposure, the half-life of the chemical in various biological compartments, and the method used to detect and measure the chemicals [58].

These data sources enable measuring exposures on a national, regional, or individual level, and at specific time points in a life course. In this paper, we provide an overview of how an exposomic framework could be applied to studies of environmental drivers of aging. We also illustrate the utility of model organisms to follow up observations made in epidemiological settings.

An ideal scenario

While a randomized clinical trial (RCT) is the gold standard design to uncover causal associations, it is unethical to conduct RCTs of exposure to many environmental toxicants. Instead, the scientific community relies on quasi-experimental observational studies. In order to apply an exposomic framework, study designs that are disease-agnostic and focused on generating a representative sample of the population of interest to enable hypothesis-free analysis of a broad range of exposure will be critical. There are several ways this could be achieved. One possibility is to create birth cohorts that are prospectively observed through their life course [59] with the following considerations:

- The cohorts should be representative of the target population of interest with representation of marginalized communities within the target population.
- Tissue samples should be collected at birth, including the meconium, placental tissue, dried blood spots, maternal blood, and cord blood, in order to assess developmental

origins of health and disease [60]. Biological samples should be collected over the course of development and adulthood. These matrices should be analyzed for: levels of environmental chemicals using high-resolution mass spectrometric techniques and characterization of genomic, epigenomic, extracellular vesicular properties, proteomic, and metabolomic status. An important variable to consider when collecting these matrices would be the unit of time at which this data should be collected [8].

- Exposure data associated with each individual in the cohort could be collected using multiple methods described earlier (Figure 2), ideally at fine-grained spatial and temporal resolution. This requires continuous updating of residence information and may also involve harmonization of exposure data across different types of monitors. Attention should also be paid to adverse childhood experiences and lived experiences, which can be collected through questionnaires [61].
- Wearable devices and sensors in the homes of participants should also be deployed to provide real-time exposure data to supplement the ecological level exposure data derived from geospatial linkages [8].

A realistic scenario

Birth cohorts are expensive, resource intensive, and prone to attrition. A complementary approach to investigate the exposome in an aging context can leverage existing infrastructure and data. Indeed, population-based cohorts with representation of different age windows are critical for an unbiased exploration of the

Exposure	Data source					
Indoor air pollution						
Pesticides						
Metals/ Metalloids						
Mutagens						
Industrial solvents						
Plasticizers and related compounds						
Ambient temperature						
Microbes						
Cigarette smoke						

Source of exposure data						
	Bio-samples		Wearable devices		Electronic health records	
Satellite remote sensing		Monitors		Questionnaires		Geo-spatial data

Figure 2. Data sources for exposure assessment. For each exposure that has been associated with hallmarks of aging listed in the first column, the corresponding source of exposure data is represented by icons along the row. Created with [BioRender.com](https://www.biorender.com)

exposome. Over decades, several cohort studies have been established that can provide the necessary data for this (Table 1). By creatively using existing data, for example, leveraging natural experiments created through changes in policies [62, 63], we can allocate resources toward the infrastructure needed to characterize the exposome, such as creating exposomic profiles of biological samples available in existing cohorts [64, 65]. Apart from integrating exposure estimates to cohort studies, it will be also be beneficial to consider ways to integrate data across cohorts representing different time points in the life course. This could be achieved by conducting parallel analyses that draw samples from cohorts at the different life course stages, like those conducted under the HELIX project in Europe [66] and in a study that leveraged data from people exposed to Arsenic in Chile and Bangladesh [67].

When using existing resources, some considerations include:

- i) Sample storage: When using banked samples, the stability of toxicants and their metabolites should be considered. Even when samples are stored in -80°C freezers, plasma-metabolite levels show storage-time-dependent changes [95].
- ii) Half-life of toxicant: Persistent environmental chemicals have long half-lives in the body whereas non-persistent chemicals are metabolized and excreted more quickly [58]. This may limit the type of chemical exposures that can be probed using banked samples that are not collected to specifically capture non-persistent exposure. By leveraging multiple biological matrices representing

different biological compartments (urine, blood, saliva, and feces), we can theoretically improve coverage of the chemical exposome.

- iii) Data harmonization: Using administrative data, residential address and history information, personal GPS data from mobile applications, and historical geospatial and satellite data, we can estimate historical and present-day exposures to several environmental factors. This data integration is no easy task. We face challenges in harmonization across different exposure estimation methods.
- iv) Privacy: It is important to also be aware of privacy concerns that may prevent study participants from sharing personal spatial data [96]. To successfully implement the use of geospatial data, researchers must convey and use methods to protect geoprivacy and disaggregate identifiable information while remaining analytically sound [97]. Further, the methods needed to access and integrate geospatial data with outcomes will differ based on population contexts [98], for example, integrating data in countries with universal health coverage will need methods different from those needed in other contexts. The readers are directed to papers by others in the field to choose an approach that protects geoprivacy [99, 100].
- v) Statistical considerations: Studying the relationship between measures that are time varying, like the exposome, and biological changes over time seen in aging poses several analytical challenges. Depending on the nature of the aging outcome, statistical methods that can be applied in these settings include: mixed-effects models [101], Cox

Table 1. A selection of population-based studies of aging well-suited to investigate the exposome

Resource	Location/cohort	Tissue type/biomarkers	Environmental data	Sample size/age range	Reference
Gateway to global aging data	US Health and Retirement Study; Mexican Health and Aging Study; English Longitudinal Study of Ageing; Study of Health, Ageing and Retirement in Europe; Costa Rican Longevity and Healthy Aging Study; Korea Employment Information Service; Japanese Study of Aging and Retirement; The Irish Longitudinal study of Ageing; The China Health and Retirement Longitudinal Study; The Longitudinal Study in India; Malaysia Ageing and Retirement Survey; Health, Aging, and Retirement in Thailand; The Brazilian Longitudinal Study of Ageing; Northern Ireland Cohort for the Longitudinal study of Ageing; Healthy Ageing in Scotland; The Health and Aging Study in Africa; Study on Global Ageing and Adult Health; Indonesia Family Life Survey	Varies by cohort. The reader is encouraged to visit this resource for information: https://gero.usc.edu/cbph/network/studies-with-bio-markers/	Residential addresses, questionnaires	The sample size varies by cohort and ranges from ~3000 to ~70 000 at baseline. The age eligibility varies by cohort and ranges from 40 to 60 years.	https://g2aging.org/
Centre for Longitudinal Studies	1958 National Child Development Study, 1970 British Cohort Study, Next Steps, Millennium Cohort Study	Blood	Questionnaires	The sample size varies by cohort but ranges from 16 000 to 19 000.	https://cls.ucl.ac.uk/
The UK Household Longitudinal Study	United Kingdom	Blood	Questionnaires	The age varies by cohort. In Wave 1, surveyed 40 000 households.	[68]
The Rotterdam Study	Netherlands	Blood	Questionnaires	Includes people of all ages. Sample size of 15 000.	[69]
The Leiden Longevity Study	Netherlands	Blood	Questionnaires	Age > 45 years. Surveyed 421 families.	[70]
The Lothian Birth Cohorts	United Kingdom	Blood	Questionnaires	Age criteria ≥89 for men and ≥91 for women. The two birth cohorts had an initial (Wave 1) sample size of 550 and 1091.	[71]
Cooperative Health Research in the Augsburg Region (KORA)	Germany	Blood	Geospatial data, questionnaires	Age > 69 years. Sample size of 18 000. Age range 25–74 years	[72]
The European Prospective Investigation into Cancer and Nutrition	Europe	Blood	Questionnaires	Sample size of 521 000. Age range 35–70 years	[73]
Longitudinal Aging Study Amsterdam, GECCO, and MINDMAP study	Netherlands	Blood	Geo-coded data, questionnaires	Initial sample size of 3805. Age range 55–84 years	[74–76]
Jerusalem Longitudinal study	Israel	Blood, serum, urine	Residential addresses, questionnaires	Initial sample size of 605. Representative of those 70-years old.	[77]
Cambridge city over 75 s cohort	United Kingdom	Blood and saliva	Residential addresses, questionnaires	Sample size 2 600. Age > 75 years	[78]

(continued)

Table 1.. (continued)

Resource	Location/cohort	Tissue type/biomarkers	Environmental data	Sample size/age range	Reference
Cognitive Function and Aging studies GAZEL cohort	United Kingdom France	Blood -	Residential address, questionnaires Occupational exposure, residential address, questionnaires	Sample size > 18 000 Age > 65 years Sample size > 20 000 Age > 35 years	[79] [80]
Heisinki Health study	Finland	-	Occupational exposure, residential address, questionnaires	Sample size > 9000 Age range 40–60 years	[81]
All of us Research Program	United States	Blood, saliva, urine	Wearable sensors, residential address, questionnaires, EHR	Aim to enroll at least 1 million people Age > 18 years	[82]
UK Biobank	United Kingdom	Blood, saliva, urine	Wearable devices, residential address, questionnaires, EHR	Sample size of 500 000 Age > 40 years	[83]
Washington Heights and Inwood Community Aging Project	United States	Blood	Residential address, questionnaires	Sample size > 4000 Age > 65 years	[84]
Multi-Ethnic Study of Atherosclerosis (MESA) and MESA Air	United States	Blood	Residential address, air pollution estimates, questionnaires	Sample size of 6814 Age range 45–84 at enrollment	[85, 86]
Mediators of Atherosclerosis in South Asians Living in America	United States	Blood, urine	Residential address, questionnaires	Sample size of 906 Age range 40–79 at enrollment	[87]
The London Life Sciences	United Kingdom	Blood	Residential address, questionnaires	Sample size ~28 000 Age range 35–74 at enrollment	[88]
Population Study Normative Aging Study	United States	Blood	Residential address, questionnaires	Initial sample size of 2280 Mean age at enrollment 42 years	[89]
Women's health initiative	United States	Blood	Residential address, questionnaires	Sample size > 161 000 Age at enrollment 50–79 years	[90]
Strong Heart Study	United States	Blood, urine	Questionnaires	Sample size > 7600 Age at enrollment 35–74 years	[91]
Baltimore Longitudinal Study of Aging	United States	Blood, urine	Residential address, questionnaires	Sample size > 3200 Age > 20 years at enrollment	[92]
Coronary Artery Risk Development in Young Adults Study	United States	Blood, urine	Residential address, questionnaires	Sample size > 5000 Age at enrollment 18–30 years	[93]
Rush Memory and Aging Project	United States	Blood	Residential address, questionnaires	Initial sample size 1556 Mean age ~81 years	[94]

Many population studies of aging exist that can be used to study environmental drivers of aging using an exposomic framework. A selection of them are shown here. GECCO, Geoscience and Health Cohort Consortium and GAZEL, GAZ and Electricite.

proportional hazards models [102], survival analyses [103], multi-state modeling techniques [104], or age-period-cohort modeling [105]. Besides handling temporal dynamics, the high-dimensionality of exposomic data needs to be treated with appropriate statistical methods [106]. Several papers have been written that describe new methods for analyzing time-varying, high-dimensional data [59].

- vi) Survival bias/mortality selection: Observations of older adults include only those individuals who have survived to advanced ages. Exposomic parameters that are more lethal may be under-represented in the oldest members of research cohorts [107]. Thus, studies that are truly representative of the population are crucial to understand real-world exposures.
- vii) Misclassification: Researchers must conduct sensitivity analyses to determine the effect of exposure misclassification or error in predictions based on sparse historical data, or other decisions and assumptions made during exposure estimation. Systematic errors that differentially affect exposure measurements in sub-groups of populations may introduce bias in analyses, for example, people with unstable residences may be assigned imprecise measures of exposure.
- viii) Study design limitations: When using a design that is a mixture of prospective and retrospective measurement, researchers would need to consider the limitations that are inherent to such a design, such as recall bias and reverse causation [36].

Toxicological follow-up

Many aspects of biological aging were first discovered in model organisms *Caenorhabditis elegans* and *Drosophila melanogaster*. Model organisms remain an important tool in uncovering mechanisms of biological aging and to discover interventions. These models can also play a role in understanding biological plausibility and gradient, which are part of Hill's causal criteria [108], of any significant findings in observational settings.

The long history of *C. elegans* in genetic and neuroscience research provides a rich resource in the pursuit of environmental drivers of aging. The large number of existing mutant libraries provides limitless opportunity and their genetic tractability makes them useful models to find as yet uncharacterized relationships between exposures and genetic susceptibility. *Caenorhabditis elegans* are amenable to high-throughput screens and have been used by the National Toxicology Program for this purpose [109]. Studies have found good correspondence between LD₅₀ values of several toxicants in rodents and *C. elegans* [110]. An example of the type of high-throughput approaches is the COPAS biosorter, which is a flow cytometer that can handle particles as large as *C. elegans*. The instrument has been used to screen for reproductive and neurodegenerative toxicants [111–113].

Over decades, researchers have developed several open source methods that favor the use of *C. elegans* in aging research. For example, the lifespan machine developed by Stroustrup et al. [114] makes it possible to measure the lifespan of thousands of worms with significantly reduced human involvement. As another example, several researchers have developed methods to study associative memory in the organism [115]. Additionally, researchers are able to characterize tissue-specific gene expression [116], miRNA expression [117], histone modifications [118], and global

metabolism [111, 119]. They are also excellent models to study reproductive aging and development [113, 120].

Data from whole organisms can be complimented by predictive toxicology, which may provide evidence of exposure, metabolism, absorption, and potential toxic effects of exposure using computational methods. The adverse outcome pathway (AOP) framework has been used to improve mechanistic understanding and to predict adverse effects of exposure using existing toxicological evidence [121]. While the use of AOP framework has generally been limited to individual chemicals, experts have recommended its use in exposome research by moving from linear pathways to networks of pathways, thus considering multiple causes for adverse effects [122]. The use of AOPs in the exposome context can help create *a priori* mechanistic links between exposure to mixtures and adverse outcomes relevant to the aging process [123].

Measures of aging

The molecular hallmarks of aging [20], established from studies of cells and model organisms, are difficult to measure in studies of humans. As a result, there remains no gold standard measure of human aging, although a range of demographic, clinical, and molecular methods have been proposed [24, 124]. In demography, researchers quantify aging at the population level from differences in mortality rates across chronological ages [125]; in clinical gerontology, researchers quantify aging at the patient level from accumulations of chronic diseases and functional deficits [126] and measures of physical frailty [127]; in the emerging field of geroscience, which seeks to translate basic science in the biology of aging to prevent chronic disease [19, 128], researchers quantify aging using algorithms that summarize omics data to estimate the state or pace of decline in system integrity, referred to as biological aging [129–131]. None of these outcomes has yet received substantial research attention in an exposomic framework. However, early studies investigating impacts of environmental pollutants suggest substantial promise [132–135].

Similar to the exposome, these processes change over time, that is, they evolve over decades. This creates the challenge of intersecting dynamic exposures to dynamic biological processes, especially since we are interested in understanding when biological aging deviates from chronological aging.

Conclusion

With a rapidly aging global population, understanding the drivers of aging is more important than ever before. The exposome framework is responsive to the reality of human exposures, they are variable in space and time, and occur concomitantly. In the paper, we have described some ways in which the exposome framework could be applied to aging research; however, challenges remain in capturing profiles of chemicals with: short half-lives, low abundance in the body, and those present in a minor subset of the population. Despite these limitations, using an exposomic framework can provide a realistic assessment of major environmental drivers of aging, providing a means to prioritize policies and interventions that can prevent unhealthy aging. Birth cohorts should be set up in several locations worldwide to ensure representation of low- and middle-income countries. International projects like the 1000 genomes project or the HAPMAP project may provide a valuable template to create exposomics profiles in populations across the globe [136, 137]. The availability of high-quality assessment of molecular hallmarks of aging is critical to studying environmental

drivers of aging and resources must be directed at improving read-outs of hallmarks of aging. Integration of historical environmental exposure data with existing population studies has the potential to accelerate this research agenda.

Funding

This work was funded by National Institutes of Health grants P30 ES009089 and UL1TR001873.

Conflict of interest statement

Dr. Miller receives royalties from his books *The Exposome: A Primer* and *The Exposome: a New Paradigm for the Environment and Health*. He also receives an annual stipend for his service as Editor-in-Chief of *Exposome*.

References

- Chen L, Deng H, Cui H, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget* 2018; 9(6):7204–7218.
- McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med*. 1998;338(3):171–179.
- Cole SW. Human social genomics. *PLOS Genet*. 2014;10(8): e1004601.
- Escher BI, Ashauer R, Dyer S, et al. Crucial role of mechanisms and modes of toxic action for understanding tissue residue toxicity and internal effect concentrations of organic chemicals. *Integr Environ Assess Manag* 2011;7(1):28–49.
- Smirnova L, Harris G, Leist M, Hartung T. Cellular resilience. *ALTEX* 2015;32(4):247–260.
- Ahima RS. Global warming threatens human thermoregulation and survival. *J Clin Invest*. 2020;130(2):559–561.
- Hoffman JB, Petriello MC, Hennig B. Impact of nutrition on pollutant toxicity: An update with new insights into epigenetic regulation. *Rev Environ Health*. 2017;32(1–2):65–72.
- Jiang C, Wang X, Li X, et al. Dynamic human environmental exposome revealed by longitudinal personal monitoring. *Cell* 2018;175(1):277–291.e31.
- Wild CP. Complementing the genome with an “exposome”: The outstanding challenge of environmental exposure measurement in molecular epidemiology. *Cancer Epidemiol Biomarkers Prev*. 2005;14(8):1847–1850.
- Wild CP. The exposome: From concept to utility. *Int J Epidemiol*. 2012;41(1):24–32.
- Niedzwiecki MM, Walker DI, Vermeulen R, Chadeau-Hyam M, Jones DP, Miller GW. The exposome: Molecules to populations. *Annu Rev Pharmacol Toxicol*. 2018. <https://doi.org/10.1146/annurev-pharmtox-010818-021315>
- Vermeulen R, Schymanski EL, Barabási A-L, Miller GW. The exposome and health: Where chemistry meets biology. *Science*. 2020;367(6476):392–396.
- Miller GW, Jones DP. The nature of nurture: Refining the definition of the exposome. *Toxicol Sci*. 2014;137(1):1–2.
- Koontz JM, Dancy BCR, Horton CL, Stallings JD, DiVito VT, Lewis JA. The role of the human microbiome in chemical toxicity. *Int J Toxicol*. 2019;38(4):251–264.
- White CM, St. John PD, Cheverie MR, Iraniparast M, Tyas SL. The role of income and occupation in the association of education with healthy aging: Results from a population-based, prospective cohort study. *BMC Public Health*. 2015;15:1181.
- Iwarsson S, Wahl H-W, Nygren C, et al. Importance of the home environment for healthy aging: Conceptual and methodological background of the European ENABLE-AGE Project. *Gerontologist*. 2007;47(1):78–84.
- Juster R-P, McEwen BS, Lupien SJ. Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neurosci Biobehav Rev*. 2010;35(1):2–16.
- Guidi J, Lucente M, Sonino N, Fava GA. Allostatic load and its impact on health: A systematic review. *Psychother Psychosom*. 2021;90(1):11–27.
- Kennedy BK, Berger SL, Brunet A, et al. Geroscience: Linking aging to chronic disease. *Cell* 2014;159(4):709–713.
- López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell* 2013;153(6):1194–1217.
- Melzer D, Pilling LC, Ferrucci L. The genetics of human ageing. *Nat Rev Genet*. 2020;21(2):88–101.
- Ruby JG, Wright KM, Rand KA, et al. Estimates of the heritability of human longevity are substantially inflated due to assortative mating. *Genetics* 2018;210(3):1109–1124.
- Rappaport SM, Smith MT. Environment and disease risks. *Science*. 2010;330(6003):460–461.
- Ferrucci L, Gonzalez-Freire M, Fabbri E, et al. Measuring biological aging in humans: A quest. *Aging Cell*. 2020;19(2). <https://doi.org/10.1111/ace1.13080>
- Whitson HE, Duan-Porter W, Schmader KE, Morey MC, Cohen HJ, Colón-Emeric CS. Physical resilience in older adults: Systematic review and development of an emerging construct. *J Gerontol A Biol Sci Med Sci*. 2016;71(4):489–495.
- Wright RO. Environment, susceptibility windows, development and child health. *Curr Opin Pediatr*. 2017;29(2):211–217.
- Terry MB, Michels KB, Brody JG, et al.; On Behalf of Breast Cancer and the Environment Research Program. Environmental exposures during windows of susceptibility for breast cancer: A framework for prevention research. *Breast Cancer Res*. 2019; 21(1):96.
- Joehanes R, Just AC, Marioni RE, et al. Epigenetic signatures of cigarette smoking. *Circ Cardiovasc Genet*. 2016;9(5):436–447.
- Saeed S, Quintin J, Kerstens HHD, et al. Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science*. 2014;345(6204):1251086.
- Saenen ND, Martens DS, Neven KY, et al. Air pollution-induced placental alterations: An interplay of oxidative stress, epigenetics, and the aging phenotype? *Clin Epigenet*. 2019;11(1):124.
- Aloizou A-M, Siokas V, Vogiatzi C, et al. Pesticides, cognitive functions and dementia: A review. *Toxicol Lett*. 2020;326: 31–51.
- Huat TJ, Camats-Perna J, Newcombe EA, Valmas N, Kitazawa M, Medeiros R. Metal toxicity links to Alzheimer's disease and neuroinflammation. *J Mol Biol*. 2019;431(9):1843–1868.
- Elkin ER, Bridges D, Loch-Carusio R. The trichloroethylene metabolite S-(1,2-dichlorovinyl)-L-cysteine induces progressive mitochondrial dysfunction in HTR-8/SVneo trophoblasts. *Toxicology* 2019;427:152283.
- Tiku V, Tan M-W, Dikic I. Mitochondrial functions in infection and immunity. *Trends Cell Biol*. 2020;30(4):263–275.
- Ivey KL, Nguyen X-MT, Posner D, et al. The structure of relationships between the human exposome and cardiometabolic health: The million veteran program. *Nutrients* 2021;13(4): 1364.
- DeBord DG, Carreón T, Lentz TJ, Middendorf PJ, Hoover MD, Schulte PA. Use of the “exposome” in the practice of epidemiology: A primer on -omic technologies. *Am J Epidemiol*. 2016; 184(4):302–314.

37. National Academies. Leveraging Advances in Remote Geospatial Technologies to Inform Precision Environmental Health Decisions—A Workshop. The National Academies Press; August 2021.
38. VoPham T, Wilson JP, Ruddell D, et al. Linking pesticides and human health: A geographic information system (GIS) and Landsat remote sensing method to estimate agricultural pesticide exposure. *Appl Geogr.* 2015;62:171–181.
39. Turner MC, Nieuwenhuijsen M, Anderson K, et al. Assessing the exposome with external measures: Commentary on the state of the science and research recommendations. *Annu Rev Public Health.* 2017;38:215–239.
40. Nuckols JR, Ward MH, Jarup L. Using geographic information systems for exposure assessment in environmental epidemiology studies. *Environ Health Perspect.* 2004;112(9):1007–1015.
41. Miranda ML, Edwards SE. Use of spatial analysis to support environmental health research and practice. *N C Med J.* 2011;72(2):132–135.
42. Sorek-Hamer M, Just AC, Kloog I. The use of satellite remote sensing in epidemiological studies. *Curr Opin Pediatr.* 2016;28(2):228–234.
43. Maxwell SK, Meliker JR, Goovaerts P. Use of land surface remotely sensed satellite and airborne data for environmental exposure assessment in cancer research. *J Expo Sci Environ Epidemiol.* 2010;20(2):176–185.
44. Carrión D, Arfer KB, Rush J, et al. A 1-km hourly air-temperature model for 13 northeastern U.S. states using remotely sensed and ground-based measurements. *Environ Res.* 2021;200:111477.
45. Solomon PA, Hopke PK, Froines J, Scheffe R. Key scientific findings and policy- and health-relevant insights from the U.S. environmental protection agency’s particulate matter supersites program and related studies: An integration and synthesis of results. *J Air Waste Manag Assoc.* 2008;58(13 Suppl):S3–92.
46. Snyder EG, Watkins TH, Solomon PA, et al. The changing paradigm of air pollution monitoring. *Environ Sci Technol.* 2013;47(20):11369–11377.
47. Hagemann R, Corsmeier U, Kottmeier C, Rinke R, Wieser A, Vogel B. Spatial variability of particle number concentrations and NO_x in the Karlsruhe (Germany) area obtained with the mobile laboratory ‘AERO-TRAM’. *Atmos Environ.* 2014;94:341–352.
48. Patel CJ, Pho N, McDuffie M, et al. A database of human exposomes and phenomes from the US National Health and Nutrition Examination Survey. *Sci Data.* 2016;3:160096.
49. Telfar Barnard LF, Baker MG, Hales S, Howden-Chapman P. Novel use of three administrative datasets to establish a cohort for environmental health research. *BMC Public Health.* 2015;15:246.
50. Namulanda G, Qualters J, Vaidyanathan A, et al. Electronic health record case studies to advance environmental public health tracking. *J Biomed Inform.* 2018;79:98–104.
51. Lin EZ, Esenther S, Mascelloni M, Irfan F, Godri Pollitt KJ. The fresh air wristband: A wearable air pollutant sampler. *Environ Sci Technol Lett.* 2020;7(5):308–314.
52. Wright SP, Hall Brown TS, Collier SR, Sandberg K. How consumer physical activity monitors could transform human physiology research. *Am J Physiol Regul Integr Comp Physiol.* 2017;312(3):R358–R367.
53. Loh M, Sarigiannis D, Gotti A, et al. How sensors might help define the external exposome. *Int J Environ Res Public Health.* 2017;14(4):434.
54. Uppal K, Walker DI, Liu K, Li S, Go Y-M, Jones DP. Computational metabolomics: A framework for the million metabolome. *Chem Res Toxicol.* 2016;29(12):1956–1975.
55. Schymanski EL, Kondić T, Neumann S, Thiessen PA, Zhang J, Bolton EE. Empowering large chemical knowledge bases for exposomics: PubChemLite meets MetFrag. *J Cheminform.* 2021;13(1):19.
56. Misra BB. New software tools, databases, and resources in metabolomics: Updates from 2020. *Metabolomics.* 2021;17(5):49.
57. Misra BB. Advances in high resolution GC–MS technology: A focus on the application of GC-Orbitrap-MS in metabolomics and exposomics for FAIR practices. *Anal Methods.* 2021;13(20):2265–2282.
58. Barr DB, Wang RY, Needham LL. Biologic monitoring of exposure to environmental chemicals throughout the life stages: Requirements and issues for consideration for the National Children’s Study. *Environ Health Perspect.* 2005;113(8):1083–1091.
59. Santos S, Maitre L, Warembourg C, et al. Applying the exposome concept in birth cohort research: A review of statistical approaches. *Eur J Epidemiol.* 2020;35(3):193–204.
60. Barker DJP. The origins of the developmental origins theory. *J Intern Med.* 2007;261(5):412–417.
61. Zota AR, VanNoy BN. Integrating intersectionality into the exposome paradigm: A novel approach to racial inequities in uterine fibroids. *Am J Public Health.* 2021;111(1):104–109.
62. Humphreys DK, Panter J, Sahlqvist S, Goodman A, Ogilvie D. Changing the environment to improve population health: A framework for considering exposure in natural experimental studies. *J Epidemiol Community Health.* 2016;70(9):941–946.
63. Crane M, Bohn-Goldbaum E, Grunseit A, Bauman A. Using natural experiments to improve public health evidence: A review of context and utility for obesity prevention. *Health Res Policy Syst.* 2020;18(1):48.
64. Terry MB, Flom J, Tehranifar P, Susser E. The role of birth cohorts in studies of adult health: the New York women’s birth cohort. *Paediatr Perinat Epidemiol.* 2009. <https://doi.org/10.1111/j.1365-3016.2009.01061.x>
65. Dennis KK, Marder E, Balshaw DM, et al. Biomonitoring in the era of the exposome. *Environ Health Perspect.* 2017;125(4):502–510.
66. Vrijheid M, Slama R, Robinson O, et al. The human early-life exposome (HELIX): Project rationale and design. *Environ Health Perspect.* 2014;122(6):535–544.
67. Bozack AK, Boileau P, Wei L, et al. Exposure to arsenic at different life-stages and DNA methylation meta-analysis in buccal cells and leukocytes. *Environ Health.* 2021;20(1):79.
68. Platt L, Knies G, Luthra R, Nandi A, Benzeval M. Understanding society at 10 years. *Eur Sociol Rev.* 2021;36(6):976–988.
69. Hofman A, Breteler MMB, van Duijn CM, et al. The Rotterdam Study: Objectives and design update. *Eur J Epidemiol.* 2007;22(11):819–829.
70. Schoenmaker M, de Craen AJM, de Meijer PHEM, et al. Evidence of genetic enrichment for exceptional survival using a family approach: The Leiden Longevity Study. *Eur J Hum Genet.* 2006;14(1):79–84.
71. Deary IJ, Gow AJ, Pattie A, Starr JM. Cohort profile: The Lothian birth cohorts of 1921 and 1936. *Int J Epidemiol.* 2012;41(6):1576–1584.
72. Holle R, Happich M, Löwel H, et al.; MONICA/KORA Study Group. KORA—A research platform for population based health research. *Gesundheitswesen.* 2005;67(Suppl 1):S19–25.

73. Riboli E, Hunt KJ, Slimani N, et al. European prospective investigation into cancer and nutrition (EPIC): Study populations and data collection. *Public Health Nutr.* 2002;5(6B):1113–1124.
74. Hoogendijk EO, Deeg DJH, de Breij S, et al. The Longitudinal Aging Study Amsterdam: Cohort update 2019 and additional data collections. *Eur J Epidemiol.* 2020;35(1):61–74.
75. Timmermans EJ, Lakerveld J, Beulens JWJ, et al. Cohort profile: The geoscience and health cohort consortium (GECCO) in the Netherlands. *BMJ Open.* 2018;8(6):e021597.
76. Beenackers MA, Doiron D, Fortier I, et al. MINDMAP: Establishing an integrated database infrastructure for research in ageing, mental well-being, and the urban environment. *BMC Public Health.* 2018;18(1):158.
77. Jacobs JM, Cohen A, Bursztyrn M, Azoulay D, Ein-Mor E, Stessman J. Cohort profile: The Jerusalem longitudinal cohort study. *Int J Epidemiol.* 2009;38(6):1464–1469.
78. Fleming J, Zhao E, O'Connor DW, Pollitt PA, Brayne C; The Cc75c Study. Cohort profile: The Cambridge City over-75s cohort (CC75C). *Int J Epidemiol.* 2007;36(1):40–46.
79. Brayne C, McCracken C, Matthews FE; Medical Research Council Cognitive Function and Ageing Study (CFAS). Cohort Profile: The medical research council cognitive function and ageing study (CFAS). *Int J Epidemiol.* 2006;35(5):1140–1145.
80. Goldberg M, Leclerc A, Zins M. Cohort profile update: The GAZEL cohort study. *Int J Epidemiol.* 2015;44(1):77–77g
81. Lahelma E, Aittomäki A, Laaksonen M, et al. Cohort profile: The Helsinki Health Study. *Int J Epidemiol.* 2013;42(3):722–730.
82. The “all of us” research program. *NE J Med* 2019;381:668–676.
83. Bycroft C, Freeman C, Petkova D, et al. The UK BioBank resource with deep phenotyping and genomic data. *Nature* 2018; 562(7726):203–209.
84. Tang MX, Cross P, Andrews H, et al. Incidence of AD in African-Americans, Caribbean Hispanics, and Caucasians in northern Manhattan. *Neurology* 2001;56(1):49–56.
85. Bild DE, Bluemke DA, Burke GL, et al. Multi-ethnic study of atherosclerosis: Objectives and design. *Am J Epidemiol.* 2002; 156(9):871–881.
86. Cohen MA, Adar SD, Allen RW, et al. Approach to estimating participant pollutant exposures in the multi-ethnic study of atherosclerosis and air pollution (MESA Air). *Environ Sci Technol.* 2009;43(13):4687–4693.
87. Kanaya AM, Kandula N, Herrington D, et al. Mediators of atherosclerosis in South Asians living in America (MASALA) study: Objectives, methods, and cohort description. *Clin Cardiol.* 2013;36(12):713–720.
88. Chambers JC, Loh M, Lehne B, et al. Epigenome-wide association of DNA methylation markers in peripheral blood from Indian Asians and Europeans with incident type 2 diabetes: A nested case-control study. *Lancet Diab Endocrinol.* 2015;3(7): 526–534.
89. Bell B, Rose CL, Damon A. The Normative Aging Study: An interdisciplinary and longitudinal study of health and aging. *Aging Hum Dev.* 1972;3(1):5–17.
90. Anderson G, Cummings S, Freedman LS, et al. Design of the women’s health initiative clinical trial and observational study. *Control Clin Trials.* 1998;19:61–109.
91. Lee ET, Welty TK, Fabsitz R, et al. The Strong Heart Study. A study of cardiovascular disease in American Indians: Design and methods. *Am J Epidemiol.* 1990;132(6):1141–1155.
92. Shock NW, Greulich RC, Andres R, et al. Normal human aging: The Baltimore Longitudinal Study of Aging (NIH Publication No. 84-2450). Bethesda, MD: National Institutes of Health; 1984.
93. Friedman GD, Cutter GR, Donahue RP, et al. CARDIA: Study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol.* 1988;41(11):1105–1116.
94. Bennett DA, Schneider JA, Buchman AS, Barnes LL, Boyle PA, Wilson RS. Overview and findings from the rush memory and aging project. *Curr Alzheimer Res.* 2012;9(6):646–663.
95. Haid M, Muschet C, Wahl S, et al. Long-term stability of human plasma metabolites during storage at -80°C . *J Proteome Res.* 2018;17(1):203–211.
96. Curtis A, Mills JW, Leitner M. Keeping an eye on privacy issues with geospatial data. *Nature* 2006;441(7090):150–150.
97. Kwan M-P, Casas I, Schmitz B. Protection of geoprivacy and accuracy of spatial information: How effective are geographical masks? *Cartographica* 2004;39(2):15–28. <https://doi.org/10.3138/X204-4223-57MK-8273>
98. Shanley L. Spatial Data Sovereignty and Privacy in Indian Country. Social Science Research Network; 2015.
99. Ajayakumar J, Curtis AJ, Curtis J. Addressing the data guardian and geospatial scientist collaborator dilemma: How to share health records for spatial analysis while maintaining patient confidentiality. *Int J Health Geogr.* 2019;18(1):30.
100. Marcon Y, Bishop T, Avraam D, et al. Orchestrating privacy-protected big data analyses of data from different resources with R and DataSHIELD. *PLoS Comput Biol.* 2021;17(3):e1008880.
101. Fitzmaurice GM, Ravichandran C. A primer in longitudinal data analysis. *Circulation* 2008;118(19):2005–2010.
102. Lamarca R, Alonso J, Gómez G, Muñoz A. Left-truncated data with age as time scale: An alternative for survival analysis in the elderly population. *J Gerontol A Biol Sci Med Sci.* 1998;53(5): M337–343. M3
103. Willett JB, Singer JD. Applications of survival analysis to aging research. *Exp Aging Res.* 1991;17(4):243–250.
104. Ruiz-Castro JE, Zenga M. A general piecewise multi-state survival model: Application to breast cancer. *Stat Methods Appl.* 2020;29(4):813–843.
105. Keyes KM, Li G. Age-period-cohort modeling. In: G Li, SP Baker, eds. *Injury Research: Theories, Methods, and Approaches.* Springer US; 2012:409–426.
106. Kalia V, Walker DI, Krasnodemski KM, Jones DP, Miller GW, Kioumourtoglou M-A. Unsupervised dimensionality reduction for exposome research. *Curr Opin Environ Sci Health.* 2020;15:32–38. <https://doi.org/10.1016/j.coesh.2020.05.001>
107. Nelson PG, Promislow DEL, Masel J. Biomarkers for aging identified in cross-sectional studies tend to be non-causative. *J Gerontol A Biol Sci Med Sci.* 2020;75(3):466–472.
108. Fedak KM, Bernal A, Capshaw ZA, Gross S. Applying the Bradford Hill criteria in the 21st century: How data integration has changed causal inference in molecular epidemiology. *Emerg Themes Epidemiol.* 2015;12(1). <https://doi.org/10.1186/s12982-015-0037-4>
109. Boyd WA, Smith MV, Co CA, et al. Developmental effects of the ToxCast™ phase I and phase II chemicals in *Caenorhabditis elegans* and corresponding responses in zebrafish, rats, and rabbits. *Environ Health Perspect.* 2016;124(5):586–593.
110. Hunt PR. The *C. elegans* model in toxicity testing. *J Appl Toxicol.* 2017;37(1):50–59.
111. Bradner JM, Kalia V, Lau FK, et al. Genetic or toxicant-induced disruption of vesicular monoamine storage and global metabolic profiling in *Caenorhabditis elegans*. *Toxicol Sci.* 2021;180(2): 313–324.
112. Pulak R. Techniques for analysis, sorting, and dispensing of *C. elegans* on the COPAS flow-sorting system. *Methods Mol Biol.* 2006;351:275–286.

113. Boyd WA, McBride SJ, Rice JR, Snyder DW, Freedman JH. A high-throughput method for assessing chemical toxicity using a *Caenorhabditis elegans* reproduction assay. *Toxicol Appl Pharmacol.* 2010;245(2):153–159.
114. Stroustrup N, Ulmschneider BE, Nash ZM, López-Moyado IF, Apfeld J, Fontana W. The *Caenorhabditis elegans* lifespan machine. *Nat Methods.* 2013;10(7):665–670.
115. Kauffman A, Parsons L, Stein G, Wills A, Kaletsky R, Murphy C. *C. elegans* positive butanone learning, short-term, and long-term associative memory assays. *J Vis Exp* 2011. <https://doi.org/10.3791/2490>
116. Kaletsky R, Yao V, Williams A, et al. Transcriptome analysis of adult *Caenorhabditis elegans* cells reveals tissue-specific gene and isoform expression. *PLoS Genet.* 2018;14(8):e1007559. <https://doi.org/10.1371/journal.pgen.1007559>
117. Martinez NJ, Ow MC, Reece-Hoyes JS, Barrasa MI, Ambros VR, Walhout AJM. Genome-scale spatiotemporal analysis of *Caenorhabditis elegans* microRNA promoter activity. *Genome Res.* 2008;18(12):2005–2015.
118. González-Aguilera C, Palladino F, Askjaer P. *C. elegans* epigenetic regulation in development and aging. *Brief Funct Genomics.* 2014;13(3):223–234.
119. Kalia V, Niedzwiecki MM, Bradner JM, et al. Cross-species metabolomic analysis of DDT and Alzheimer's disease-associated tau toxicity. *bioRxiv* 2021; 2021.06.14.448355.
120. Hunt PR, Olejnik N, Bailey KD, Vaught CA, Sprando RL. *C. elegans* development and activity test detects mammalian developmental neurotoxins. *Food Chem Toxicol.* 2018;121:583–592.
121. Ankley GT, Bennett RS, Erickson RJ, et al. Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk assessment. *Environ Toxicol Chem.* 2010; 29(3):730–741.
122. Knapen D, Vergauwen L, Villeneuve DL, Ankley GT. The potential of AOP networks for reproductive and developmental toxicity assay development. *Reprod Toxicol.* 2015;56:52–55.
123. Escher BI, Hackermüller J, Polte T, et al. From the exposome to mechanistic understanding of chemical-induced adverse effects. *Environ Int.* 2017;99:97–106.
124. Hägg S, Belsky DW, Cohen AA. Developments in molecular epidemiology of aging. *Emerg Top Life Sci.* 2019;3(4):411–421.
125. Baudisch A. The pace and shape of ageing. *Methods Ecol Evol.* 2011;2(4):375–382.
126. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci.* 2007;62(7):722–727.
127. Fried LP, Tangen CM, Walston J, et al.; Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: Evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001; 56(3):M146–156.
128. Campisi J, Kapahi P, Lithgow GJ, Melov S, Newman JC, Verdin E. From discoveries in ageing research to therapeutics for healthy ageing. *Nature* 2019;571(7764):183–192.
129. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of ageing. *Nat Rev Genet.* 2018;19(6):371–384.
130. Belsky DW, Caspi A, Arseneault L, et al. Quantification of the pace of biological aging in humans through a blood test, the DunedinPoAm DNA methylation algorithm. *eLife.* 2020;9:e54870.
131. Lehallier B, Gate D, Schaum N, et al. Undulating changes in human plasma proteome profiles across the lifespan. *Nat Med.* 2019;25(12):1843–1850.
132. Ward-Caviness CK, Nwanaji-Enwerem JC, Wolf K, et al. Long-term exposure to air pollution is associated with biological aging. *Oncotarget* 2016;7(46):74510–74525.
133. Ward-Caviness CK, Russell AG, Weaver AM, et al. Accelerated epigenetic age as a biomarker of cardiovascular sensitivity to traffic-related air pollution. *Aging (Albany NY).* 2020;12(23):24141–24155.
134. White AJ, Kresovich JK, Keller JP, et al. Air pollution, particulate matter composition and methylation-based biologic age. *Environ Int.* 2019;132:105071.
135. Schmidt CW. Environmental factors in successful aging: The potential impact of air pollution. *Environ Health Perspect.* 2019;127(10):102001.
136. Auton A, Brooks LD, Durbin RM, et al.; 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 2015;526(7571):68–74.
137. Frazer KA, Ballinger DG, Cox DR, et al.; International HapMap Consortium. A second generation human haplotype map of over 3.1 million SNPs. *Nature* 2007;449(7164):851–861.